

ON THE FROBENIUS-SCHUR INDICATORS FOR QUASI-HOPF ALGEBRAS

PETER SCHAUENBURG

ABSTRACT. Mason and Ng have given a generalization to semisimple quasi-Hopf algebras of Linchenko and Montgomery's generalization to semisimple Hopf algebras of the classical Frobenius-Schur theorem for group representations. We give a simplified proof, in particular a somewhat conceptual derivation of the appropriate form of the Frobenius-Schur indicator that indicates if and in which of two possible fashions a given simple module is self-dual.

1. INTRODUCTION

Let (throughout this paper) k be an algebraically closed field of characteristic zero.

Let G be a finite group, and V a finite-dimensional k -linear representation of G . According to the classical Frobenius-Schur Theorem, V falls into one of the following three mutually exclusive classes: Either it is not selfdual, or there is a nondegenerate G -invariant bilinear form on V that is symmetric, respectively skew-symmetric. Which of the three cases is at hand can be decided by computing the Frobenius-Schur indicator $\nu_V = \chi_V(|G|^{-1} \sum_{g \in G} g^2)$. This can only take the values $0, 1, -1$, which occur when V is not selfdual, admits a nondegenerate symmetric invariant bilinear form, or admits a nondegenerate skew-symmetric invariant bilinear form, respectively.

Linchenko and Montgomery [10] have generalized the Frobenius-Schur Theorem to irreducible representations V of a semisimple Hopf algebra H , which are found to fall into the same three classes, again detected by the Frobenius-Schur indicator, now given by $\nu_V = \chi_V(\Lambda_{(1)}\Lambda_{(2)})$ for a normalized integral $\Lambda \in H$.

Finally, Mason and Ng [11] have generalized the results of Linchenko and Montgomery to the case of a semisimple quasi-Hopf algebra H . As pointed out by Mason and Ng (and also by Pavel Etingof in an email to Susan Montgomery), a quite general argument essentially due to Fuchs, Ganchev, Szlachanyi, and Vescerny s [4] shows that a trichotomy generalizing that in the Frobenius-Schur and Linchenko-Montgomery Theorems can be obtained for simple objects of any pivotal fusion category — this covers the representation category of a semisimple quasi-Hopf algebra by results of Etingof, Nikshych, and Ostrik [3]. It is perhaps more surprising that a formula for a Frobenius-Schur indicator detecting the three cases can again be found, by adorning the formula from [10] with the structure elements of a quasi-Hopf algebra in a rather non-obvious fashion.

1991 *Mathematics Subject Classification.* 16W30.

Key words and phrases. Quasi-Hopf algebra, Frobenius-Schur indicator.

The author is supported by a Heisenberg-Stipendium from the DFG (Deutsche Forschungsgemeinschaft).

The present paper contains no new results in this direction. We do, however, present a new proof for the fact that the Frobenius-Schur indicator of Mason and Ng does in fact indicate what it should. We find this worthwhile because our proof does not involve a lot of calculations with the structure elements and axioms of quasi-Hopf algebras, and is quite short. We should acknowledge that on our short and direct route to the indicator formula we do not encounter (and prove) many of the facts contained in the work of Mason and Ng.

Key to our proof is writing the Frobenius-Schur indicator of a module V as the trace of a certain endomorphism of the invariant subspace of $V \otimes V$. In the case of ordinary Hopf algebras, where the endomorphism is given by flipping tensor factors, this argument appears in work of Kashina, Sommerhäuser, and Zhu [8].

2. DEFINITIONS AND PRELIMINARIES

A monoidal category $\mathcal{C}$ is a category equipped with a bifunctor $\otimes: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$, a coherent natural (in $U, V, W \in \mathcal{C}$ isomorphism $\Phi: (U \otimes V) \otimes W \rightarrow U \otimes (V \otimes W)$), and a neutral object I , which, for simplicity, we assume strict in the sense that $I \otimes V = V = V \otimes I$, naturally in $V \in \mathcal{C}$. Coherence of Φ (and the neutral object) means that all reasonable diagrams involving only instances of Φ and introducing and cancelling neutral objects commute. A consequence is that $\mathcal{C}$ is equivalent as a monoidal category to a strict monoidal category, that is, one in which Φ is an identity. We refer to [9] for background on monoidal categories.

Except for a few facts on general monoidal categories, we will mostly deal with vector spaces over a fixed algebraically closed field k of characteristic zero, and $\otimes$ will mean tensor product over k . We let $\tau: V \otimes W \rightarrow W \otimes V$ denote the flip of tensor factors. We will recklessly use the formal notation $v \otimes w \in V \otimes W$ for general elements of $V \otimes W$, even if we know that the element in question is not a simple tensor. In particular, notations like $q = q^{(1)} \otimes q^{(2)} \in V \otimes W$ will often be used to refer to the components of a tensor.

A quasibialgebra H is an associative algebra (whose multiplication map we denote by $\nabla: H \otimes H \rightarrow H$) with an algebra map $\Delta: H \rightarrow H \otimes H$ called comultiplication, an algebra map $\varepsilon: H \rightarrow k$ that is a counit for Δ , and an invertible element $\phi \in H \otimes H \otimes H$ such that the category ${}_H\mathcal{M}$ is a monoidal category as follows: The tensor product of $V, W \in {}_H\mathcal{M}$ is taken over k and endowed with the diagonal module structure $h(v \otimes w) = h_{(1)}v \otimes h_{(2)}w$, where $\Delta(h) = h_{(1)} \otimes h_{(2)}$, the coherent associator isomorphism $\Phi: (U \otimes V) \otimes W \rightarrow U \otimes (V \otimes W)$ for $U, V, W \in {}_H\mathcal{M}$ is left multiplication by ϕ . We write $\phi = \phi^{(1)} \otimes \phi^{(2)} \otimes \phi^{(3)}$ and $\phi^{-1} = \phi^{(-1)} \otimes \phi^{(-2)} \otimes \phi^{(-3)}$ when in need of referring to the tensor components of ϕ . The neutral object in ${}_H\mathcal{M}$ is the trivial module k with action via ε , and the usual canonical identifications $k \otimes V = V = V \otimes k$ for $V \in {}_H\mathcal{M}$.

We made a point of not actually recalling the axioms on the data Δ and ϕ that ensure that ${}_H\mathcal{M}$ is monoidal — we will never have to use most of those axioms. We will need to know that applying ε to any of the three tensor factors of ϕ yields $1 \in H \otimes H$. For details and background on quasibialgebras and quasi-Hopf algebras, which were introduced by Drinfeld [2], we again refer to [9].

A quasiantipode (S, α, β) for a quasibialgebra H consists of an anti-algebra endomorphism S of H , and elements $\alpha, \beta \in H$, such that

$$\begin{aligned} S(h_{(1)})\alpha h_{(2)} &= \varepsilon(h)\alpha, & h_{(1)}\beta S(h_{(2)}) &= \varepsilon(h)\beta, \\ \phi^{(1)}\beta S(\phi^{(2)})\alpha\phi^{(3)} &= 1, & S(\phi^{(-1)})\alpha\phi^{(-2)}\beta\phi^{(-3)} &= 1 \end{aligned}$$

hold in H , for $h \in H$. A quasi-Hopf algebra is a quasibialgebra with a quasiantipode. If H is finite-dimensional then S is always a bijection by a result of Bulacu and Caenepeel [1]. If H has a quasiantipode, then the category of finite dimensional H -modules is rigid: Every finite-dimensional module $V \in {}_H\mathcal{M}$ has a dual object $V^\vee \in {}_H\mathcal{M}$, whose underlying vector space is the vector space dual V^* , endowed with the module structure in which $h \in H$ acts as the dual of the map given by the action of $S(h)$ on V . The evaluation and coevaluation (or “dual basis”) morphisms are given by

$$\begin{aligned} \text{ev}: V^* \otimes V &\ni \varphi \otimes v \mapsto \varphi(\alpha v) \in k \\ \text{db}: k &\ni 1 \mapsto \beta v_i \otimes v^i. \end{aligned}$$

where $v_i \otimes v^i \in V \otimes V^*$ is the canonical element.

If H has bijective antipode, then the functor $(-)^\vee$ is an equivalence. We denote the inverse functor by ${}^\vee(-)$ and call ${}^\vee V$ the right dual of V . Just like the left dual, the right dual has V^* as its underlying vector space.

As for any object V of a monoidal category having a dual object $V^\vee$, we obtain an adjunction

$$A_0: \text{Hom}_H(U, V \otimes W) \cong \text{Hom}_H(V^\vee \otimes U, W)$$

by

$$\begin{aligned} A_0(f) &= \left(V^\vee \otimes U \xrightarrow{V^\vee \otimes f} V^\vee \otimes (V \otimes W) \xrightarrow{\Phi^{-1}} (V^\vee \otimes V) \otimes W \xrightarrow{\text{ev} \otimes W} W \right) \\ A_0^{-1}(g) &= \left(U \xrightarrow{\text{db} \otimes U} (V \otimes V^\vee) \otimes U \xrightarrow{\Phi} V \otimes (V^\vee \otimes U) \xrightarrow{V \otimes g} V \otimes W \right) \end{aligned}$$

We recall the following elements of $H \otimes H$ introduced by Hausser and Nill [5, 6]:

$$\begin{aligned} q_R &= \phi^{(1)} \otimes S^{-1}(\alpha\phi^{(3)})\phi^{(2)} & p_R &= \phi^{(-1)} \otimes \phi^{(-2)}\beta S(\phi^{(-3)}) \\ q_L &= S(\phi^{(-1)})\alpha\phi^{(-2)} \otimes \phi^{(-3)} & p_L &= \phi^{(2)}S^{-1}(\phi^{(1)}\beta) \otimes \phi^{(3)} \end{aligned}$$

Writing $f(u) = f^{(1)}(u) \otimes f^{(2)}(u) \in V \otimes W$ for $f: U \rightarrow V \otimes W$, we find

$$A_0(f)(\varphi \otimes u) = (\phi^{(-1)}\varphi)(\alpha\phi^{(-2)}f^{(1)}(u))\phi^{(-3)}f^{(2)}(u) = \varphi(q_L^{(1)}f^{(-1)}(u))q_L^{(2)}f^{(-2)}(u)$$

and

$$A_0^{-1}(g)(u) = \phi^{(1)}\beta v_i g(\phi^{(2)}v^i \otimes \phi^{(3)}u) = v_i g(p_L^{(1)}v^i \otimes p_L^{(2)}u).$$

Specializing $U = k$ and using the identification $\text{Hom}_H(k, M) = M^H = \{m \in M \mid hm = \varepsilon(h)m \forall h \in H\}$, we find

$$(2.1) \quad A: (V \otimes W)^H \xrightarrow{\sim} \text{Hom}_H(V^\vee, W); \quad A(v \otimes w)(\varphi) = \varphi(q_L^{(1)}v)q_L^{(2)}w$$

with $A^{-1}(g) = \beta v_i g(v^i)$. We note $v_i \otimes A(v \otimes w)(v^i) = q_L(v \otimes w)$.

Finally, recall [7] that a quasi-Hopf algebra H is semisimple if and only if it contains a normalized integral Λ , that is, an element $\Lambda \in H$ satisfying $h\Lambda = \varepsilon(h)\Lambda$ for all $h \in H$ and $\varepsilon(\Lambda) = 1$. For any H -module M , the H -linear projection $\pi: M \rightarrow M^H$ onto the isotypical component of the trivial module, or the invariant

subspace $M^H = \{m \in M \mid hm = \varepsilon(h)m \forall h \in H\}$, is multiplication by the normalized integral Λ .

In the sequel, we will refer to all the structure elements mentioned for a (semisimple) quasi-Hopf algebra in this section without further notice.

3. THE PIVOTAL STRUCTURE

A pivotal structure for a monoidal category $\mathcal{C}$ with a duality functor $(-)^{\vee}$ is by definition a monoidal natural isomorphism $j: V \rightarrow V^{\vee\vee}$. We note that (as a special case of a fact pertaining to all monoidal transformations) we have $(j_V)^{\vee} = j_{V^{\vee}}^{-1}$ for all $V \in \mathcal{C}$.

Recall [3] that for a morphism $f: V \rightarrow V^{\vee\vee}$ in a rigid monoidal category one can define the (categorical) trace of f by

$$\text{tr}_V(f) = \left(I \xrightarrow{\text{db}} V \otimes V^{\vee} \xrightarrow{f \otimes V^{\vee}} V^{\vee\vee} \otimes V^{\vee} \xrightarrow{\text{ev}} I \right) \in \text{End}_{\mathcal{C}}(V)$$

We will depend heavily upon the following result of Etingof, Nikshych, and Ostrik: By [3, Prop.8.24 and Prop.8.23], the monoidal category ${}_H\mathcal{M}$ for a semisimple quasi-Hopf algebra is pivotal. More precisely, for any simple H -module V there is a unique isomorphism $j_V: V \rightarrow V^{\vee\vee}$ with $\text{tr}_V(j_V) = \dim V$. The j_V are the components at the simple modules of a unique natural transformation j , which is a pivotal structure on ${}_H\mathcal{M}$, and satisfies $\text{tr}_V(j_V) = \dim V$ for all finite-dimensional H -modules V .

Since j is natural, there exists a unique invertible $g \in H$ with $j(v) = \iota(gv)$ for all $v \in V$, where $\iota: V \rightarrow V^{**}$ is the canonical vector space isomorphism.

Since the invariant subspace $(V \otimes W)^H$ is isomorphic to $\text{Hom}_H(k, V \otimes W)$, we see that $V^{\vee} \otimes V$ contains a unique up to scaling H -invariant element for every simple V . The formula for the categorical trace of $j_{\vee V}$ shows that the unique invariant element in $V^{\vee} \otimes V$ mapped to $\dim V$ under the evaluation map $V^{\vee} \otimes V \rightarrow k$ is $(j_{\vee V} \otimes V) \text{db}_{\vee V}(1) = j(\beta v^i) \otimes v_i = \beta j(v^i) \otimes v_i = \beta v^i \otimes j^{-1}(\iota(v_i)) = \beta v^i \otimes g^{-1}v_i$.

On the other hand, it is easy to exhibit elements in $V^{\vee} \otimes V$ mapped to $\dim V$ under the evaluation map: For $t = t^{(1)} \otimes t^{(2)} \in H \otimes H$, the element $t^{(1)}v^i \otimes t^{(2)}v_i$ is mapped to $(t^{(1)}v^i)(\alpha t^{(2)}v_i) = \chi_V(S(t^{(1)})\alpha t^{(2)})$. For this to be equal to $\dim V = \chi_V(1)$ it suffices to have $S(t^{(1)})\alpha t^{(2)} = 1$, which is the case both for $t = p_L$ and $t = p_R$. Since the evaluation morphism is H -linear, we have $\text{ev}(\Lambda\xi) = \text{ev}(\xi)$ for every $\xi \in V^{\vee} \otimes V$. If $\text{ev}(\xi) = \dim V$, then it follows that $\Lambda\xi = \beta v^i \otimes g^{-1}v_i$. We conclude

Lemma 3.1. *Let H be a semisimple quasi-Hopf algebra and $t \in \{p_L, p_R\} \subset H \otimes H$. For every finite-dimensional H -module V we have*

$$\beta v^i \otimes g^{-1}v_i = \Lambda_{(1)}t^{(1)}v^i \otimes \Lambda_{(2)}t^{(2)}v_i \in V^{\vee} \otimes V$$

and $g^{-1}S(\beta) = S(\Lambda_{(1)}t^{(1)})\Lambda_{(2)}t^{(2)} \in H$.

Proof. We proved the first formula for the case where V is simple. This entails that the second formula holds in $\text{End}_k(V)$ for every simple module, hence in H , which in turn implies the first formula for any module. $\square$

Remark 3.2. The second formula gives an explicit formula for g in the case that β is a unit in H , and comes close otherwise. However, the first, somewhat less explicit formula proves particularly useful in our computations below.

4. FROBENIUS-SCHUR INDICATORS

Let V and W be two objects of a pivotal rigid monoidal category $\mathcal{C}$. Then we have a bijection

$$T = T_{VW}: \mathcal{C}(V^\vee, W) \rightarrow \mathcal{C}(W^\vee, V),$$

given by $T(f) = (W^\vee \xrightarrow{f^\vee} V^{\vee\vee} \xrightarrow{j_V^{-1}} V)$. We note that

$$T_{WV}T_{VW}(f) = j_W^{-1}(j_V^{-1}f^\vee)^\vee = j_W^{-1}f^{\vee\vee}(j_V^{-1})^\vee = fj_V^{-1}(j_V^{-1})^\vee = f$$

In particular, T_{VV} is an order two automorphism of $\mathcal{C}(V^\vee, V)$. In the case that V is a self-dual simple module in $\mathcal{C} = {}_H\mathcal{M}$, T_{VV} is multiplication by a scalar thanks to Schur's lemma, and the scalar can only be ± 1 . Thus, there are exactly three possibilities for a simple H -module V : Either V is not self-dual, in which case we put $\mu_V = 0$, or it is, and every isomorphism $f: V^\vee \rightarrow V$ satisfies $j_V f = \mu_V f^\vee$, with the sign $\mu_V \in \{\pm 1\}$ independent of the choice of f .

Linchenko and Montgomery [10], generalizing a classical result on group representations by Frobenius and Schur, have proved the above trichotomy for simple modules of a semisimple Hopf algebra (note j is trivial in this case by the Larson-Radford Theorem), and shown that which of the cases applies to a particular module can be decided using the Frobenius-Schur indicator, given by the formula $\nu_V(V) = \chi_V(\Lambda_{(1)}\Lambda_{(2)})$. The indicator can only take the values $0, \pm 1$, depending on which of the three possible behaviors V displays.

Mason and Ng [11] have in turn generalized the results of Linchenko and Montgomery to the quasi-Hopf algebra case. We have already seen that the same trichotomy is preserved here. It can be reformulated as in [11] to be a property of bilinear forms on V : An isomorphism $f: V \rightarrow V^*$ is the same as a nondegenerate bilinear form $\sigma: V \otimes V \rightarrow k$, by the correspondence $\sigma(v, w) = f(v)(w)$. One checks that f is an H -linear map $f: V \rightarrow V^\vee$ if and only if $\sigma(hv, w) = \sigma(v, S(h)w)$, that is, σ is a bilinear form with adjoint S in the sense of [11]. If f corresponds to σ , then $f^\vee j_V$ corresponds to σ' given by $\sigma'(v, w) = \sigma(w, gv)$. Thus, the trichotomy can be restated as follows: If there exists a nondegenerate bilinear form σ on V with adjoint S , then there is $\mu_V \in \{\pm 1\}$ so that $\sigma(v, w) = \mu_V \sigma(w, gv)$ for any such form. If there is no such form, we put $\mu_V = 0$ again.

In the rest of this section we will deal with the question how one can come up with a modification of the Frobenius-Schur indicator of Linchenko and Montgomery suitable for the quasi-Hopf algebra case. This means that, rather than only label the three possible relations of V to its dual by a number μ_V , we are looking for a formula computing a number $\nu_V \in k$ from the structure elements of a semisimple quasi-Hopf algebra and the character χ_V of a simple module V , such that $\nu_V = \mu_V$ (and in particular $\nu_V \in \{0, \pm 1\}$).

We begin by considering two H -modules V, W and computing the composition

$$E_{VW} := \left((V \otimes W)^H \xrightarrow{A} \text{Hom}_H(V^\vee, W) \xrightarrow{T_{V,W}} \text{Hom}_H(W^\vee, V) \xrightarrow{A^{-1}} (W \otimes V)^H \right),$$

where A was introduced in (2.1). We have

$$\begin{aligned}
A^{-1}TA(v \otimes w) &= \beta w_i \otimes TA(v \otimes w)(w^i) \\
&= \beta w_i \otimes j_V^{-1}A(v \otimes w)^\vee(w^i) \\
&= \beta A(v \otimes w)(v^i) \otimes j_V^{-1}(\iota(v_i)) \\
&= A(v \otimes w)(\beta v^i) \otimes g^{-1}(v_i) \\
&= A(v \otimes w)(\Lambda_{(1)}t^{(1)}v^i) \otimes \Lambda_{(2)}t^{(2)}v_i \\
&= \Lambda_{(1)}t^{(1)}A(v \otimes w)(v^i) \otimes \Lambda_{(2)}t^{(2)}v_i \\
&= \Lambda_{(1)}t^{(1)}q_L^{(2)}w \otimes \Lambda_{(2)}t^{(2)}q_L^{(1)}v
\end{aligned}$$

for $v \otimes w \in (V \otimes W)^H$, so that

$$E_{VW} = \left((V \otimes W)^H \hookrightarrow V \otimes W \xrightarrow{q_L} V \otimes W \xrightarrow{\tau} W \otimes V \xrightarrow{t} W \otimes V \xrightarrow{\pi} (W \otimes V)^H \right).$$

Now consider the case where $V = W$ is simple. Then $(V \otimes V)^H$ is zero unless V is selfdual, and then $(V \otimes V)^H$ is one-dimensional. In either case T_{VV} is multiplication by $\mu_V \in \{0, \pm 1\}$. Consequently E_{VV} is multiplication by the same scalar μ_V , and this is also the trace of E_{VV} . But E_{VV} has the same trace as

$$\begin{aligned}
E' &:= \left(V \otimes V \xrightarrow{\pi} (V \otimes V)^H \hookrightarrow V \otimes V \xrightarrow{q_L} V \otimes V \xrightarrow{\tau} V \otimes V \xrightarrow{t} V \otimes V \right) \\
&= \left(V \otimes V \xrightarrow{q_L \Delta(\Lambda)} V \otimes V \xrightarrow{\tau} V \otimes V \xrightarrow{t} V \otimes V \right).
\end{aligned}$$

Now using the general observation

$$(4.1) \quad \text{tr}((f_1 \otimes f_2) \circ \tau \circ (f_3 \otimes f_4)) = \text{tr}(f_3 f_1 f_4 f_2)$$

for endomorphisms $f_1, \dots, f_4$ of V we find that

$$\begin{aligned}
\text{tr}(E_{VV}) &= \text{tr}(E') \\
&= \chi_V(q_L^{(1)}\Lambda_{(1)}t^{(1)}q_L^{(2)}\Lambda_{(2)}t^{(2)}) \\
&= \chi_V(\nabla(q_L \Delta(\Lambda)t)).
\end{aligned}$$

We have proved the following result of Mason and Ng:

Theorem 4.1. *Let H be a semisimple quasi-Hopf algebra, and $t \in \{p_L, p_R\}$. Put $\nu_H := \nabla(q_L \Delta(\Lambda)t)$. Let V be a simple H -module, and define the Frobenius-Schur indicator $\nu_V := \chi_V(\nu_H)$. Then $\nu_V \in \{0, \pm 1\}$, and the following are equivalent for $\mu \in \{\pm 1\}$:*

- (1) $\nu_V = \mu$
- (2) *There is an H -linear isomorphism $f: V \rightarrow V^\vee$ with $f^\vee j_V = \mu f$.*
- (3) *There is a nondegenerate bilinear form σ on V with adjoint S and $\sigma(v, w) = \mu \sigma(w, gv)$ for all $v, w \in V$.*

In closing we should stress that the key trick of writing the indicator as the trace of an endomorphism of $(V \otimes V)^H$, using (4.1), is due to Kashina, Sommerhäuser, and Zhu [8]; they treat the case of ordinary Hopf algebras, where E_{VW} is just the flip of tensor factors.

REFERENCES

- [1] BULACU, D., AND CAENEPEEL, S. Integrals for (dual) quasi-Hopf algebras. applications. *preprint* (math.QA/0110063).
- [2] DRINFEL'D, V. G. Quasi-Hopf algebras. *Leningrad Math. J.* 1 (1990), 1419–1457.
- [3] ETINGOF, P., NIKSHYCH, D., AND OSTRIK, V. On fusion categories. *preprint* (math.QA/0203060).
- [4] FUCHS, J., GANCHEV, C., SZLACHÁNYI, K., AND VESCERNYÉS, P. S_4 -symmetry of $6j$ -symbols and Frobenius-Schur indicators in rigid monoidal C^* -categories. *J. Math. Phys.* 40 (1999), 408–426.
- [5] HAUSSER, F., AND NILL, F. Diagonal crossed products by duals of quasi-quantum groups. *Rev. Math. Phys.* 11 (1999), 553–629.
- [6] HAUSSER, F., AND NILL, F. Doubles of quasi-quantum groups. *Comm. Math. Phys.* 199 (1999), 547–589.
- [7] HAUSSER, F., AND NILL, F. Integral theory for quasi-Hopf algebras. *preprint* (math.QA/9904164).
- [8] KASHINA, Y., SOMMERHÄUSER, Y., AND ZHU, Y. On higher Frobenius-Schur indicators.
- [9] KASSEL, C. *Quantum Groups*, vol. 155 of *GTM*. Springer, 1995.
- [10] LINCHENKO, V., AND MONTGOMERY, S. A Frobenius-Schur theorem for Hopf algebras.
- [11] MASON, G., AND NG, S.-H. Central invariants and Frobenius-Schur indicators for semisimple quasi-Hopf algebras. *preprint* (math.QA/0303213).

DEPARTMENT OF MATHEMATICS, USC, LOS ANGELES, CA 90089-1113, USA, AND MATHEMATISCHES INSTITUT DER UNIVERSITÄT MÜNCHEN, THERESIENSTR. 39, 80333 MÜNCHEN, GERMANY
E-mail address: `schauen@mathematik.uni-muenchen.de`